



# Soil Facts

## *Poultry Manure as a Fertilizer Source*

Poultry manure is an excellent source of nutrients and can be incorporated into most fertilizer programs. Those using manures must practice sound soil fertility management to prevent nutrient imbalances and associated animal health risks, as well as surface-water and groundwater contamination. The key to successful management is to match the nutritional requirements of the crop with nutrients available in the manure. The value of poultry manure varies not only with its nutrient composition and availability, but also with management and handling costs.

### Nutrient Composition and Sampling Procedure

The nutrient composition of poultry manure varies with the type of bird, the feed ration, the proportion of litter to droppings, the manure handling system, and the type of litter. Consequently, all manures should be sampled and analyzed for specific nutrient content before you apply them to the land. Waste samples can be analyzed for \$4 by the North Carolina Department of Agriculture (NCDA), Agronomic Division, Plant Analysis Lab, P.O. Box 27647, Blue Ridge Road Center, Raleigh, NC 27611. Other qualified private laboratories can also perform the analysis (fees vary).

Collecting a representative manure sample

is essential to reliable nutrient analysis. The nutrient value of litter varies greatly within the poultry house. To reduce sample variability, collect subsamples of broiler, turkey, and duck litter in 6 to 12 areas of the house. Samples taken around waterers, feeders, and brooders should be proportionate to the space these areas occupy in the house. At each location, collect litter by digging an area down to the earth; be careful, however, not to include soil. Place the subsamples in a plastic bucket, mix thoroughly, and put 2 to 3 pounds of the mixture in a sample container. Samples from stockpiled litter should be taken from at least 6 locations around the pile, all at depths of at least 18 inches. Subsamples should be

**Table 1. Average Nutrient Composition of Broiler Manures**

Manure Type	Total N	Ammonium NH <sub>4</sub>	Phosphorus P <sub>2</sub> O <sub>5</sub>	Potassium K <sub>2</sub> O
Fresh (no litter)	26	10	17	11
Broiler house litter <sup>1</sup>	72	11	78	46
Roaster house litter <sup>1</sup>	73	12	75	45
Breeder house litter <sup>1</sup>	31	7	54	31
Stockpiled litter <sup>1</sup>	36	8	80	34

<sup>1</sup>Annual manure and litter accumulation; typical litter base is sawdust, wood shavings, or peanut hulls.

Source: *Biological and Agricultural Engineering Department, NCSU.*

Distributed in furtherance of the Acts of Congress of May 8 and June 30, 1914. Employment and program opportunities are offered to all people regardless of race, color, national origin, sex, age, or disability. North Carolina State University, North Carolina A&T State University, U.S. Department of Agriculture, and local governments cooperating.



**North Carolina Cooperative Extension Service**

NORTH CAROLINA STATE UNIVERSITY  
COLLEGE OF AGRICULTURE & LIFE SCIENCES

mixed and submitted as suggested for litter from poultry houses.

To increase sample uniformity in poultry manure slurries and lagoon sludges, stir them before sampling. Within an anaerobic lagoon, liquids are relatively uniform above the sludge zone; nevertheless, take several subsamples and combine them.

If you cannot have the manure analyzed, use the mean nutrient values for your specific type of poultry manure found in Tables 1 through 4. Table 5 gives the average values for the secondary and micro-nutrients ordinarily listed in the manure analysis report.

When using mean values for manure nutrient composition, exercise caution to avoid over- or under-fertilization. Also, after several years, elements such as copper or zinc may accumulate and reach very high levels. To avoid these problems, take an annual plant tissue and a biennial soil sample to monitor nutrient levels.

## Nutrient Availabilities

Except for nitrogen, the availability of most nutrients in poultry manures is fairly consistent. Nitrogen can occur in several forms, each of which can be lost when subjected to different management or environmental conditions.

Nitrogen in poultry wastes comes from uric acid, ammonia salts, and organic (fecal) matter. The predominant form is uric acid, which readily transforms to ammonia ( $\text{NH}_3$ ), a gaseous form of nitrogen that can evaporate if not mixed into the soil. When it is thoroughly mixed, the ammonia changes to ammonium ( $\text{NH}_4^+$ ), which can be temporarily held on clay particles and organic matter. Thus, soil mixing can reduce nitrogen losses and increase the amount available to plants.

Table 6 lists the first-year nutrient availability coefficients for

**Table 2. Average Nutrient Composition of Layer Manures**

Manure Type	Total N	Ammonium $\text{NH}_4\text{-N}$	Phosphorus $\text{P}_2\text{O}_5$	Potassium $\text{K}_2\text{O}$
Fresh (no litter)	26	6	22	11
Undercage scraped <sup>1</sup>	28	14	31	20
Highrise stored <sup>2</sup>	38	18	56	30
lb/1,000 gallons				
Liquid slurry <sup>3</sup>	62	42	59	37
Anaerobic lagoon sludge	26	8	92	13
lb/acre-inch				
Anaerobic lagoon liquid	179	154	46	266

<sup>1</sup>Manure collected within two days.

<sup>2</sup>Annual manure accumulation on unpaved surfaces.

<sup>3</sup>Six-12 months' accumulation of manure, excess water usage, and storage-surface rainfall surplus; does not include fresh water for flushing.

Source: Biological and Agricultural Engineering Department, NCSU.

**Table 3. Average Nutrient Composition of Turkey Manures**

Manure Type	Total N	Ammonium $\text{NH}_4\text{-N}$	Phosphorus $\text{P}_2\text{O}_5$	Potassium $\text{K}_2\text{O}$
Fresh (no litter)	27	8	25	12
Brooder house litter <sup>1</sup>	45	9	52	32
Grower house litter <sup>2</sup>	57	16	72	40
Stockpiled litter <sup>3</sup>	36	8	72	33

<sup>1</sup>Based on cleanout after each flock.

<sup>2</sup>Based on annual cleanout after full production.

<sup>3</sup>Based on annual house accumulation removed to uncovered stockpile to be spread within six months.

Source: Biological and Agricultural Engineering Department, NCSU.

**Table 4. Average Nutrient Composition of Duck Manures**

Manure Type	Total N	Ammonium $\text{NH}_4\text{-N}$	Phosphorus $\text{P}_2\text{O}_5$	Potassium $\text{K}_2\text{O}$
Fresh (no litter)	28	5	23	17
House litter <sup>1</sup>	19	3	17	14
Stockpiled litter <sup>2</sup>	24	5	42	22

<sup>1</sup>Annual manure and litter accumulation; typical litter base is wood shavings.

<sup>2</sup>Annual house accumulation removed to uncovered stockpile to be spread within six months.

Source: Biological and Agricultural Engineering Department, NCSU.

**Table 5. Average Secondary and Micronutrient Content of Poultry Manures**

Manure Type	Ca	Mg	S	Na	Fe	Mn	B	Mo	Zn	Cu
lb/ton										
<b>Layer</b>										
Undercage scraped	43.0	6.1	7.1	4.5	0.52	0.27	0.050	0.00390	0.32	0.036
Highrise stored	86.0	6.0	8.8	5.0	1.8	0.52	0.046	0.00038	0.37	0.043
<b>Broiler Litter</b>										
Broiler house	41.0	8.0	15.0	13.0	1.3	0.67	0.054	0.00085	0.63	0.45
Roaster house	43.0	8.5	14.0	13.0	1.6	0.74	0.049	0.00082	0.68	0.51
Breeder house	94.0	6.8	8.5	8.6	1.3	0.57	0.035	0.00048	0.52	0.21
Stockpiled	54.0	8.0	12.0	6.2	1.5	0.59	0.041	0.00069	0.55	0.27
<b>Turkey Litter</b>										
Brooder house	28.0	5.7	7.6	5.9	1.4	0.52	0.047	0.00081	0.46	0.36
Grower house	42.0	7.0	10.0	8.4	1.3	0.65	0.048	0.00092	0.64	0.51
Stockpiled	42.0	6.8	9.5	6.4	1.5	0.62	0.047	0.00095	0.56	0.34
<b>Duck Litter</b>										
Duck house	22.0	2.7	3.1	2.8	.98	0.31	0.021	0.00040	0.26	.056
Stockpiled	27.0	4.4	5.6	8.8	1.2	0.47	0.030	0.00030	0.47	.050
lb/1,000 gallons										
<b>Layer</b>										
Liquid slurry	35.0	6.8	8.2	5.3	2.9	0.42	0.040	0.018	0.43	0.080
Lagoon sludge	71.0	7.2	12.0	4.2	2.2	2.3	0.082	0.014	0.80	0.14
lb/acre-inch										
<b>Layer</b>										
Lagoon liquid	25.0	7.4	52.0	51.0	2.0	0.24	0.37	0.020	0.70	0.19

various poultry manures. Determine the available nutrients by multiplying these values by the nutrient composition values listed on the waste analysis report or in Tables 1 through 4. The NCDA's Agronomic Division calculates available nutrients and lists them in its report.

## Application Rate

Land application rates are generally determined by matching the available nitrogen or phosphorus content of the waste to the nutrient requirements of the crop. In most cases, nitrogen requirements determine the application rate, unless the area is designated "nutrient sensitive" and

indicates that phosphorus movement off-site can lead to eutrophication of surface waters. In nondesignated areas, phosphorus movement can be adequately controlled with conservation measures such as grass field borders, grassed waterways, contour planting, and reduced tillage, which minimize soil and residual manure movement. Leaching of phosphorus is extremely limited on mineral soils and should not contribute to ground-water contamination.

Nitrogen recommendations for various crops are listed in Table 7. Use these rates as guidelines with realistic yield capabilities for the crop and field. With feed and forage crops, excessive manure application

can produce high nitrate concentrations, which can harm livestock (through nitrate poisoning) and promote nutrient imbalances that may lead to grass tetany. If loading rates are to be based on phosphorus, apply the amount suggested by soil-test recommendations. Because the manure may not supply adequate amounts of all the other nutrients required by the crop, be sure to take a soil test and, if necessary, supplement with commercial fertilizer.

In addition to monitoring nutrients, be sure to maintain an adequate soil pH, which will help to maximize crop yields and nutrient availability

and promote the decomposition of organic matter. The biological conversion of organic matter to nitrate is an acid-forming process. Take annual or biennial soil samples to monitor pH changes. When livestock wastes are applied at agronomic rates, high salinity (excess salt) has not been a problem, given normal amounts of rainfall in North Carolina.

A worksheet to help you determine land application rates is included at the end of the text.

## Timing and Uniformity of Manure Applications

To minimize nitrogen losses, apply manure as near as possible to planting time or to the crop growth stage during which nitrogen is most needed. Surfacewater and groundwater contaminations are greater in areas of high rainfall and when manures for spring crops are applied in fall or winter. For coarse-textured soils, manures should be applied frequently and at low rates throughout the growing season because such soils have a high water infiltration rate and a low ability to hold nutrients. Unused nitrogen can therefore be lost by leaching.

Exercise caution when applying lagoon liquid by irrigation on crops undergoing stress (for example, corn during an extended drought). A heavy coating of manure solids on the leafy vegetation can cause ammonia burn. Except in extreme cases, this damage is usually short term and does not significantly reduce yields. With concentrated lagoon liquids, use several small applications rather than one large dose.

Whether poultry waste is applied by manure spreaders or irrigation systems, you *must* apply it uniformly. A lack of uniformity leads to nutrient excesses and deficiencies, lower yields, and variable crop moisture at harvest time.

**Table 6. First-Year Nitrogen Availability Coefficients for Different Poultry Manures**

Manure Type	Soil			
	Injection <sup>1</sup>	Incorporation <sup>2</sup>	Broadcast <sup>3</sup>	Irrigation <sup>4</sup>
P <sub>2</sub> O <sub>5</sub> and K <sub>2</sub> O availability coefficients				
All manure types	0.8	0.8	0.7	0.7
N availability coefficient				
All poultry litters <sup>5</sup>	—	0.6	0.5	—
Layers (no litter)	—	0.6	0.4	—
Layer anaerobic lagoon sludge	0.6	0.6	0.4	0.4
Layer anaerobic liquid slurry	0.8	0.7	0.4	0.3
Layer liquid lagoon	0.9	0.8	0.5	0.5

<sup>1</sup>Manure injected directly into soil and covered immediately.

<sup>2</sup>Surface-spread manure plowed or disked into soil within two days.

<sup>3</sup>Surface-spread manure uncovered for one month or longer.

<sup>4</sup>Sprinkler-irrigated liquid uncovered for one month or longer.

<sup>5</sup>Includes in-house and stockpiled litters.

**Table 7. Nitrogen Fertilization Guidelines**

Commodity	lb N/RYE <sup>1</sup>
Corn (grain)	1.0 - 1.25 lb N/bu
Corn (silage)	10 - 20 lb N/ton
Cotton	0.06 - 0.12 lb N/lb lint
Sorghum (grain)	2.0 - 2.5 lb N/cwt
Wheat (grain)	1.7 - 2.4 lb N/bu
Rye (grain)	1.7 - 2.4 lb N/bu
Barley (grain)	1.4 - 1.6 lb N/bu
Triticale (grain)	1.4 - 1.6 lb N/bu
Oats	1.0 - 1.3 lb N/bu
Bermudagrass (hay <sup>2,3</sup> )	40 - 50 lb N/dry ton
Tall fescue (hay <sup>2,3</sup> )	40 - 50 lb N/dry ton
Orchardgrass (hay <sup>2,3</sup> )	40 - 50 lb N/dry ton
Small grain(hay <sup>2,3</sup> )	50 - 60 lb N/dry ton
Sorghum-sudangrass (hay <sup>2,3</sup> )	45 - 55 lb N/dry ton
Millet (hay <sup>2,3</sup> )	45 - 55 lb N/dry ton
Pine and hardwood trees <sup>4</sup>	40 60 lb N/acre/year

<sup>1</sup>RYE = Realistic Yield Expectation

<sup>2</sup>Annual maintenance guidelines

<sup>3</sup>Reduce N rate by 25 percent when grazing.

<sup>4</sup>On trees less than 5 feet tall, N will stimulate undergrowth competition.

**Table 8. Minimum Amount of Land Needed to Apply Poultry Manure as a Nitrogen Fertilizer Source (Based on the Nitrogen Rate Required by the Crop)**

Manure Handling and Production Unit	Soil Incorporated <sup>1</sup>				Surface Broadcast <sup>2</sup>			
	lb N/acre/year							
	100	200	300	400	100	200	300	400
	<b>Annual acres/1,000 bird single capacity</b>							
<b>Layer</b>								
Undercaged scraped	4.80	2.40	1.60	1.20	3.00	1.50	1.00	0.75
Highrise scraped	4.30	2.15	1.43	1.07	2.60	1.30	0.87	0.65
Liquid manure/slurry	6.70	3.35	2.23	1.68	4.00	2.00	1.33	1.00
Anaerobic lagoon sludge	0.71	0.35	0.24	0.18	0.56	0.28	0.19	0.14
Anaerobic lagoon liquid	0.87	0.43	0.29	0.22	0.84	0.42	0.28	0.21
<b>Broiler Litter</b>								
Broiler house	2.40	1.20	0.80	0.60	1.96	0.98	0.65	0.49
Roaster house	4.30	2.15	1.43	1.08	3.60	1.80	1.20	0.90
Breeder house	4.70	2.35	1.57	1.18	3.20	1.60	1.07	0.80
Stockpiled	1.20	0.60	0.40	0.30	0.92	0.46	0.31	0.23
<b>Turkey Litter</b>								
Poult house	1.40	0.70	0.47	0.35	1.08	0.54	0.36	0.27
Brooder house	8.10	4.05	2.70	2.02	5.60	2.80	1.87	1.40
Grower hen house	5.70	2.85	1.90	1.43	4.00	2.00	1.33	1.00
Grower tom house	8.60	4.30	2.87	2.15	6.00	3.00	2.00	1.50
<b>Stockpiled</b>								
Poult	0.94	0.47	0.31	0.23	0.76	0.38	0.25	0.19
Hen	3.00	1.50	1.00	0.75	2.40	1.20	0.80	0.60
Tom	4.50	2.25	1.50	1.13	3.60	1.80	1.20	0.90
<b>Duck Litter</b>								
Duck house	3.00	1.50	1.00	0.75	2.20	1.10	0.73	0.55
Stockpiled	1.50	0.75	0.50	0.38	1.08	0.54	0.36	0.27

<sup>1</sup>Incorporated within two days

<sup>2</sup>Not incorporated for at least 1 month

### Acreage Requirements for New Facilities

Whenever manure or lagoon liquid samples are available for analysis, they should be used to determine application rates and acreage requirements. When you are planning new facilities, however, the average values can help determine approximate acreage requirements for a poultry operation of a given size.

Table 8 gives minimum acreage requirements for various nitrogen fertilization rates. This table can be used to estimate the minimum acreage required to use all of the manure.

Suppose that a producer is interested in building two broiler houses with a combined 50,000 bird capacity/growout. The producer is planning to spread this litter on a bermudagrass hay field capable of

producing 6 dry tons per acre. From Table 7, the bermudagrass will require 300 lb nitrogen per acre (6 tons x 50 lb N/dry ton). How many acres of bermudagrass would be needed for the entire year's waste? Using Table 8, under surface broadcast column 300, we find that each 1,000-bird capacity would require 0.65 acres for land application of broiler litter. For a 50,000-bird growout operation (0.65 x 50), the

producer would need 32.5 acres for a year's worth of litter.

## Value of Manure

When comparing manure to commercial fertilizers, convert total manure nutrients to available nutrients by using the availability coefficients. Consider the following example. Analysis of the available nitrogen, phosphorus ( $P_2O_5$ ), and potassium ( $K_2O$ ) content in a broiler litter sample that will be incorporated shows that it contains 43 pounds of nitrogen per ton, 62 pounds of phosphate per ton, and 37 pounds of potash per ton. The current fertilizer prices for nitrogen, phosphate, and potash are as follows: \$0.23 per pound of nitrogen; \$0.22 per pound of phosphate; and \$0.12 per pound of potash as potassium chloride. One ton of broiler litter would be worth the following:

$$(43 \times \$0.23) + (62 \times \$0.22) + (37 \times \$0.12) = \$27.97 \text{ per ton}$$

This value does not cover hauling, handling, or application costs, nor does it include the value of other essential nutrients available in the manure. In addition, it assumes that the soil test has recommended each nutrient, when, in fact, many may not be needed. Nutrients not needed should not be considered when you assess the manure's value.

## Land Application Worksheet

Farmer Jones is preparing to spread broiler litter on a field and incorporate it within two days to supply nutrients to his corn crop. Last year, he grew soybeans in the field.

His corn-yield goal is 140 bushels per acre, and he has decided to apply the equivalent of 140 pounds of nitrogen per acre (Table 7). His land is not subject to erosion, is not in a nutrient-sensitive watershed, and has grassed borders and waterways to further reduce the potential of

**Table 9. Estimated Residual Nitrogen Provided by Legumes Grown in Rotation**

Legume <sup>1</sup>	Residual Nitrogen Available
	lb/acre
Alfalfa <sup>2</sup>	80 - 100
Harry vetch <sup>2</sup>	80 - 100
Crimson clover <sup>2</sup>	60 - 75
Austrian winter pea <sup>2</sup>	50 - 60
Soybeans <sup>3</sup> harvested for seed	15 - 30
Peanuts <sup>3</sup> harvested for seed	20 - 40

<sup>1</sup>Assumes good stand.

<sup>2</sup>Killed before planting current spring crop.

<sup>3</sup>Legume planted in previous year or season. More nitrogen available if fall-planted crop immediately follows legume; less nitrogen available with spring-planted crop.



**Figure 1. One way to apply poultry manure is to use a litter spreader.**

runoff.

Farmer Jones used a starter fertilizer on his corn crop at a rate to supply 10 pounds of nitrogen per acre and 34 pounds of phosphorus per acre. He intends to supply the rest of his nitrogen needs by applying broiler litter with a litter spreader (Figure 1) and incorporating it within two days.

How much litter does he need to spread in order to meet the nitrogen needs of his corn crop? Will he need

to supplement the crop with additional potash or phosphate to satisfy his soil-test recommendations of 50 pounds per acre of each nutrient? The answers are given in the following worksheet. Use Table 9 to estimate available nitrogen carry-over from legumes.

**Worksheet: Determining the Nutrient Needs of Your Crop**

	Example	Your Farm
1. Crop to be grown	corn	_____
2. Total nutrients required		
a. N (Table 7) (lb/acre)	140	_____
b. P <sub>2</sub> O <sub>5</sub> (soil test) (lb/acre)	50	_____
c. K <sub>2</sub> O (soil test) (lb/acre)	50	_____
3. Pounds of starter or preplant fertilizer used		
a. N (lb/acre)	10	_____
b. P <sub>2</sub> O <sub>5</sub> (lb/acre)	34	_____
c. K <sub>2</sub> O (lb/acre)	0	_____
4. Residual N credit from legumes (Table 9) (lb/acre)	20	_____
5. Net nutrient needs of crop (lb/acre)		
Nitrogen: total need (item 2a) minus additional N from starter (item 3a) minus legume residual (item 4)		
a. N: 140 – 10 – 20 (lb/acre)	110	_____
Phosphorus and potassium: total need (items 2b and 2c) minus additional nutrients from starter (items 3b and 3c)		
b. P <sub>2</sub> O <sub>5</sub> : 50 – 34 (lb/acre)	16	_____
c. K <sub>2</sub> O: 50 – 0 (lb/acre)	50	_____
6. Nutrient totals in manure. If analysis report already gives available nutrients, skip this item.		
a. Total N (Tables 1-4 or waste samples) (lb/ton)	72	_____
b. P <sub>2</sub> O <sub>5</sub> (lb/ton)	78	_____
c. K <sub>2</sub> O (lb/ton)	46	_____
7. Nutrients available to crop (items 6a, 6b, and 6c) times availability coefficients (Table 6). If analysis report already gives available nutrients, fill in those numbers.		
a. Available N: 72 x 0.6 (lb/ton)	43.2	_____
b. Available P <sub>2</sub> O <sub>5</sub> : 78 x 0.8 (lb/ton)	62.4	_____
c. Available K <sub>2</sub> O: 46 x 0.8 (lb/ton)	36.8	_____
8. Application rate to supply priority nutrient		
a. Priority nutrient	nitrogen	_____
b. Amount of priority nutrient needed (lb/acre from item 5a)	110	_____
c. Rate of manure needed to supply priority nutrient (item 8b) divided by (item 7a): 110/43.2 (tons/acre)	2.55	_____

**Worksheet (continued)**

**Rate of Manure to Apply**

	Example	Your Farm
9. Pounds per acre of all nutrients supplied at the application rate required to meet the needs for the priority nutrient: for each nutrient, enter the available nutrients (items 7a, 7b, and 7c) times manure rate (item 8c)		
a. N supplied: $43.2 \times 2.55$ (lb/acre)	<u>110</u>	_____
b. $P_2O_5$ supplied: $62.4 \times 2.55$ (lb/acre)	<u>159</u>	_____
c. $K_2O$ supplied: $36.8 \times 2.55$ (lb/acre)	<u>94</u>	_____
10. Nutrient balance: net nutrient need (-) or excess (+) after application of manure at calculated rate. Subtract the net nutrient needs of the crop (items 5a, 5b, and 5c) from the nutrient rate applied (items 9a, 9b, and 9c).		
a. N balance: $110 - 110$ (lb/acre)	<u>0</u>	_____
b. $P_2O_5$ balance: $159 - 16$ (lb/acre)	<u>+143</u>	_____
c. $K_2O$ balance: $92 - 50$ (lb/acre)	<u>+44</u>	_____

Note: Calculation format modified from Pennsylvania Department of Environmental Resources, *Field Application of Manure*, October 1986.

*Prepared by*

*J. P. Zublena, Extension Soil Science Specialist;  
J. C. Barker, Extension Agricultural Engineering Specialist; and  
T. A. Carter, Extension Poultry Science Specialist.*

*The authors wish to acknowledge the assistance and cooperation of the North Carolina Department of Agriculture's Agronomic Division in the analysis of samples and the development of the data used in this publication.*

*10,000 copies of the public document were printed at a cost of \$1,698.00, or \$.17 per copy.*

Published by  
**NORTH CAROLINA COOPERATIVE EXTENSION SERVICE**